



Mapping multiple QTL in experimental crosses

Karl W Broman

Department of Biostatistics
Johns Hopkins University

www.biostat.jhsph.edu/~kbroman


Human vs mouse




www.daviddeen.com



Backcross




Intercross



5

Genetic map




7

Phenotype data


Sugiyama et al. *Genomics* 71:70-77, 2001

250 male mice from the backcross (A × B) × B
 Blood pressure after two weeks drinking water with 1% NaCl



6

Genotype data




8

Goals

- Identify quantitative trait loci (QTL) (and interactions among QTL)
- Interval estimates of QTL location
- Estimated QTL effects

9

Statistical structure



The missing data problem:

Markers \longleftrightarrow QTL


The model selection problem:

QTL, covariates \longrightarrow phenotype

10

ANOVA at marker loci

- Split mice into groups according to genotype at a marker.
- Do a t-test / ANOVA.
- Repeat for each marker.



11

Interval mapping

Lander & Botstein (1989)

- Assume a single QTL model.
- Consider each position in the genome, one at a time, as the location of the putative QTL.
- Let $q = 0/1$ if the (unobserved) QTL genotype is BB/AB. (Or $0/1/2$ if the QTL genotype is AA/AB/BB in an intercross.)

Assume $y \mid q \sim N(\mu_q, \sigma)$

- Calculate $p_q = \Pr(q \mid \text{marker data})$.

$y \mid \text{marker data} \sim \sum_q p_q \phi(y \mid \mu_q, \sigma)$

12

LOD scores


$\text{LOD}(\lambda) = \log_{10}$ likelihood ratio comparing the hypothesis of a QTL at position λ versus that of no QTL

$$= \log_{10} \left\{ \frac{\Pr(y|\text{QTL at } \lambda, \hat{\mu}_{q\lambda}, \hat{\sigma}_\lambda)}{\Pr(y|\text{no QTL}, \hat{\mu}, \hat{\sigma})} \right\}$$

$\hat{\mu}_{q\lambda}, \hat{\sigma}_\lambda$ are the MLEs, assuming a single QTL at position λ .

No QTL model: The phenotypes are iid $N(\mu, \sigma^2)$.


Permutation test



13


15

LOD curves




14

Permutation results




16

LOD curves




17

LOD curves




17

LOD curves



17

Estimated effects



18

Modeling multiple QTL

- Reduce residual variation → increased power
- Separate linked QTL
- Identify interactions among QTL (epistasis)

19

2-dim, 2-QTL scan

For each pair of positions, fit the following models:

$$H_f : y = \mu + \beta_1 q_1 + \beta_2 q_2 + \gamma q_1 q_2 + \epsilon$$

$$H_a : y = \mu + \beta_1 q_1 + \beta_2 q_2 + \epsilon$$

$$H_1 : y = \mu + \beta_1 q_1 + \epsilon$$

$$H_0 : y = \mu + \epsilon$$

$$\text{LOD}_f(s, t) = \log_{10} \{ L_f(s, t) / L_0 \}$$

20

2-dim, 2-QTL scan

For each pair of positions, fit the following models:

$$H_f : y = \mu + \beta_1 q_1 + \beta_2 q_2 + \gamma q_1 q_2 + \epsilon$$

$$H_a : y = \mu + \beta_1 q_1 + \beta_2 q_2 + \epsilon$$

$$H_1 : y = \mu + \beta_1 q_1 + \epsilon$$

$$H_0 : y = \mu + \epsilon$$

20

2-dim, 2-QTL scan

For each pair of positions, fit the following models:

$$H_f : y = \mu + \beta_1 q_1 + \beta_2 q_2 + \gamma q_1 q_2 + \epsilon$$

$$H_a : y = \mu + \beta_1 q_1 + \beta_2 q_2 + \epsilon$$


$$H_1 : y = \mu + \beta_1 q_1 + \epsilon$$

$$H_0 : y = \mu + \epsilon$$

$$\text{LOD}_i(s, t) = \log_{10} \{ L_f(s, t) / L_a(s, t) \}$$


20

LOD_i and LOD_f




21

LOD_i and LOD_{fv1}




22

LOD_i and LOD_f




21

Estimated effects



23

Chr 1: LOD_i and LOD_{av1}



24

Hypothesis testing?

- In the past, QTL mapping has been regarded as a task of hypothesis testing.

Is this a QTL?

Much of the focus has been on adjusting for test multiplicity.

- It is better to view the problem as one of model selection.

What set of QTL are well supported?

Is there evidence for QTL-QTL interactions?

Model = a defined set of QTL and QTL-QTL interactions (and possibly covariates and QTL-covariate interactions).

25

Model selection

- Class of models
 - Additive models
 - + pairwise interactions
 - + higher-order interactions
 - Regression trees
- Model fit
 - Maximum likelihood
 - Haley-Knott regression
 - extended Haley-Knott
 - Multiple imputation
 - MCMC
- Model comparison
 - Estimated prediction error
 - AIC, BIC, penalized likelihood
 - Bayes
- Model search
 - Forward selection
 - Backward elimination
 - Stepwise selection
 - Randomized algorithms

26

Target

- Selection of a model includes two types of errors:
 - Miss important terms (QTLs or interactions)
 - Include extraneous terms
- Unlike in hypothesis testing, we can make both errors at the same time.
- Identify as many correct terms as possible, while controlling the rate of inclusion of extraneous terms.

27

What is special here?

- Goal: identify the major players
- A continuum of ordinal-valued covariates (the genetic loci)
- Association among the covariates
 - Loci on different chromosomes are independent
 - Along chromosome, a very simple (and known) correlation structure

28

Exploratory methods

- Condition on a large-effect QTL
 - Reduce residual variation
 - Conditional LOD score:

$$\text{LOD}(q_2 | q_1) = \log_{10} \left\{ \frac{\text{Pr}(\text{data} | q_1, q_2)}{\text{Pr}(\text{data} | q_1)} \right\}$$

- Piece together the putative QTL from the 1d and 2d scans
 - Omit loci that no longer look interesting (drop-one-at-a-time analysis)
 - Study potential interactions among the identified loci
 - Scan for additional loci (perhaps allowing interactions), conditional on these

29

Automation

- Assistance to the masses
- Understanding performance
- Many phenotypes

30

Additive QTL

Simple situation:

- Dense markers
- Complete genotype data
- No epistasis

$$y = \mu + \sum \beta_j q_j + \epsilon \quad \text{which } \beta_j \neq 0?$$

$$\text{BIC}(\gamma) = \log \text{RSS}(\gamma) + \left(\frac{\log n}{n} \right) |\gamma|$$

31

Additive QTL

Simple situation:

- Dense markers
- Complete genotype data
- No epistasis

$$y = \mu + \sum \beta_j q_j + \epsilon \quad \text{which } \beta_j \neq 0?$$

$$\text{BIC}_\delta(\gamma) = \log \text{RSS}(\gamma) + \left(\delta \cdot \frac{\log n}{n} \right) |\gamma|$$

31

Additive QTL

Simple situation:

- Dense markers
- Complete genotype data
- No epistasis

$$y = \mu + \sum \beta_j q_j + \epsilon \quad \text{which } \beta_j \neq 0?$$

$$\text{LOD}_\delta(\gamma) = \text{LOD}(\gamma) - (2\delta \log n) |\gamma|$$

31

Additive QTL

Simple situation:

- Dense markers
- Complete genotype data
- No epistasis

$$y = \mu + \sum \beta_j q_j + \epsilon \quad \text{which } \beta_j \neq 0?$$

$$\text{LOD}_\delta(\gamma) = \text{LOD}(\gamma) - T |\gamma|$$

31

Additive QTL

Simple situation:

- Dense markers
- Complete genotype data
- No epistasis

$$y = \mu + \sum \beta_j q_j + \epsilon \quad \text{which } \beta_j \neq 0?$$

$$\text{LOD}_\delta(\gamma) = \text{LOD}(\gamma) - T |\gamma|$$

0 vs 1 QTL: $\text{LOD}_\delta(\emptyset) = 0$

$$\text{LOD}_\delta(\{\lambda\}) = \text{LOD}(\{\lambda\}) - T$$

31

Additive QTL

Simple situation:

- Dense markers
- Complete genotype data
- No epistasis

$$y = \mu + \sum \beta_j q_j + \epsilon \quad \text{which } \beta_j \neq 0?$$

$$\text{LOD}_\delta(\gamma) = \text{LOD}(\gamma) - T |\gamma|$$

For the mouse genome:

$$T = 2.69 \text{ (BC) or } 3.52 \text{ (F}_2\text{)}$$

31

Experience

- Controls rate of inclusion of extraneous terms
- Forward selection over-selects
- Forward selection followed by backward elimination works as well as MCMC
- Need to define performance criteria
- Need large-scale simulations

Broman & Speed, JRSS B 64:641-656, 2002

32

Epistasis

$$y = \mu + \sum \beta_j q_j + \sum \gamma_{jk} q_j q_k + \epsilon$$

$$\text{LOD}_{\delta\epsilon}(\gamma) = \text{LOD}(\gamma) - T_m |\gamma|_m + T_i |\gamma|_i$$

T_m = as chosen previously

T_i = ?

33

Idea 1

Imagine there are two additive QTL and consider a 2d, 2-QTL scan.

$$T_i = 95\text{th percentile of the distribution of} \\ \max \text{LOD}_f(s, t) - \max \text{LOD}_a(s, t)$$

34

Idea 1

Imagine there are two additive QTL and consider a 2d, 2-QTL scan.

$$T_i = 95\text{th percentile of the distribution of} \\ \max \text{LOD}_f(s, t) - \max \text{LOD}_a(s, t)$$

For the mouse genome:

$$T_m = 2.69 \text{ (BC) or } 3.52 \text{ (F}_2\text{)}$$

$$T_i^H = 2.62 \text{ (BC) or } 4.28 \text{ (F}_2\text{)}$$

34

Idea 2

Imagine there is one QTL and consider a 2d, 2-QTL scan.

$$T_m + T_i = 95\text{th percentile of the distribution of} \\ \max \text{LOD}_f(s, t) - \max \text{LOD}_1(s)$$

For the mouse genome:

$$T_m = 2.69 \text{ (BC) or } 3.52 \text{ (F}_2\text{)}$$

$$T_i^H = 2.62 \text{ (BC) or } 4.28 \text{ (F}_2\text{)}$$

$$T_i^L = 1.19 \text{ (BC) or } 2.69 \text{ (F}_2\text{)}$$

35


Idea 2

Imagine there is one QTL and consider a 2d, 2-QTL scan.

$$T_m + T_i = 95\text{th percentile of the distribution of} \\ \max \text{LOD}_f(s, t) - \max \text{LOD}_1(s)$$


35

Models as graphs




36

Results



37


Results



$$T_m = 2.69 \quad T_i^H = 2.62 \quad T_i^L = 1.19 \quad T_m + T_i^H = 5.31 \quad T_m + T_i^L = 3.88$$


37

Results



37


Add an interaction?



$$T_m = 2.69 \quad T_i^H = 2.62 \quad T_i^L = 1.19 \quad T_m + T_i^H = 5.31 \quad T_m + T_i^L = 3.88$$

38


Add an interaction?



$$T_m = 2.69 \quad T_i^H = 2.62 \quad T_i^L = 1.19 \quad T_m + T_i^H = 5.31 \quad T_m + T_i^L = 3.88$$

38


Add an interaction?



$$T_m = 2.69 \quad T_i^H = 2.62 \quad T_i^L = 1.19 \quad T_m + T_i^H = 5.31 \quad T_m + T_i^L = 3.88$$

38


Add an interaction?



$$T_m = 2.69 \quad T_i^H = 2.62 \quad T_i^L = 1.19 \quad T_m + T_i^H = 5.31 \quad T_m + T_i^L = 3.88$$

38


Add an interaction?



$$T_m = 2.69 \quad T_i^H = 2.62 \quad T_i^L = 1.19 \quad T_m + T_i^H = 5.31 \quad T_m + T_i^L = 3.88$$

38


Add another QTL?



$$T_m = 2.69 \quad T_i^H = 2.62 \quad T_i^L = 1.19 \quad T_m + T_i^H = 5.31 \quad T_m + T_i^L = 3.88$$

39


Add another QTL?



$$T_m = 2.69 \quad T_i^H = 2.62 \quad T_i^L = 1.19 \quad T_m + T_i^H = 5.31 \quad T_m + T_i^L = 3.88$$

39


Add another QTL?



$$T_m = 2.69 \quad T_i^H = 2.62 \quad T_i^L = 1.19 \quad T_m + T_i^H = 5.31 \quad T_m + T_i^L = 3.88$$

39

Add a pair of QTL?



$$T_m = 2.69 \quad T_i^H = 2.62 \quad T_i^L = 1.19 \quad T_m + T_i^H = 5.31 \quad T_m + T_i^L = 3.88$$

39

To do

- Improve search procedures
- Study performance
(especially relative to other approaches)
- Measuring model uncertainty
- Measuring uncertainty in QTL location

40

Summary

- QTL mapping is a model selection problem
- The criterion for comparing models is most important
- We're focusing on a penalized likelihood method and are close to a practiceable solution

42

Bayes/MCMC

Advantages

- All analysis aspects combined
- More fully captures uncertainty
- More clean expression of uncertainty in the inference

Disadvantages

- May require a specialist
- Prior specification is difficult
- Bayes factors can be difficult to interpret
- Can be difficult to assess performance

41

Acknowledgments

Ani Manichaikul	Johns Hopkins University
Gary Churchill	Jackson Laboratory
Šaunak Sen	University of California, San Francisco
Terry Speed	University of California, Berkeley
Brian Yandell	University of Wisconsin, Madison
Fumihiko Sugiyama	now at University of Tsukuba, Japan
Bev Paigen	Jackson Laboratory

43