

Mapping multiple QTL in experimental crosses

Karl W Broman

Department of Biostatistics and Medical Informatics
University of Wisconsin – Madison

www.biostat.wisc.edu/~kbroman


[→ Teaching → Miscellaneous lectures]

Example


Sugiyama et al. Genomics 71:70-77, 2001

250 male mice from the backcross $(A \times B) \times B$

Blood pressure after two weeks drinking water with 1% NaCl




Genetic map



3

Genotype data




4

Goals

- Identify quantitative trait loci (QTL)
(and interactions among QTL)
- Interval estimates of QTL location
- Estimated QTL effects


5

LOD curves



6

Estimated effects



7

Modeling multiple QTL

- Reduce residual variation → increased power
- Separate linked QTL
- Identify interactions among QTL (epistasis)

8

2-dim, 2-QTL scan

For all pairs of positions, fit the following models:

$$H_f : y = \mu + \beta_1 q_1 + \beta_2 q_2 + \gamma q_1 q_2 + \epsilon$$

$$H_a : y = \mu + \beta_1 q_1 + \beta_2 q_2 + \epsilon$$

$$H_1 : y = \mu + \beta_1 q_1 + \epsilon$$

$$H_0 : y = \mu + \epsilon$$

\log_{10} likelihoods:

$$l_f(s, t) \quad l_a(s, t) \quad l_1(s) \quad l_0$$

9

2-dim, 2-QTL scan

LOD scores:


$$\text{LOD}_f(s, t) = l_f(s, t) - l_0$$

$$\text{LOD}_a(s, t) = l_a(s, t) - l_0$$

$$\text{LOD}_i(s, t) = l_f(s, t) - l_a(s, t)$$


$$\text{LOD}_1(s) = l_1(s) - l_0$$

Results: LOD_i and LOD_f



11

Results: LOD_i and LOD_f



12

Summaries

Consider each pair of chromosomes, (j, k) ,
and let $c(s)$ denote the chromosome for position s .

$$M_f(j, k) = \max_{c(s)=j, c(t)=k} \text{LOD}_f(s, t)$$

$$M_a(j, k) = \max_{c(s)=j, c(t)=k} \text{LOD}_a(s, t)$$

$$M_1(j, k) = \max_{c(s)=j \text{ or } k} \text{LOD}_1(s)$$


$$M_i(j, k) = M_f(j, k) - M_a(j, k)$$

$$M_{fv1}(j, k) = M_f(j, k) - M_1(j, k)$$

$$M_{av1}(j, k) = M_a(j, k) - M_1(j, k)$$


13

Results: LOD_i and LOD_{fv1}




14

Estimated effects



15

Chr 1: LOD_i and LOD_{av1}



16

Hypothesis testing?

- In the past, QTL mapping has been regarded as a task of hypothesis testing.

Is this a QTL?

Much of the focus has been on adjusting for test multiplicity.

- It is better to view the problem as one of model selection.

What set of QTL are well supported?

Is there evidence for QTL-QTL interactions?

Model = a defined set of QTL and QTL-QTL interactions
(and possibly covariates and QTL-covariate interactions).

17

Model selection

- Class of models
 - Additive models
 - + pairwise interactions
 - + higher-order interactions
 - Regression trees
- Model comparison
 - Estimated prediction error
 - AIC, BIC, penalized likelihood
 - Bayes
- Model fit
 - Maximum likelihood
 - Haley-Knott regression
 - extended Haley-Knott
 - Multiple imputation
 - MCMC
- Model search
 - Forward selection
 - Backward elimination
 - Stepwise selection
 - Randomized algorithms

18

Target

- Selection of a model includes two types of errors:
 - Miss important terms (QTLs or interactions)
 - Include extraneous terms
- Unlike in hypothesis testing, we can make both errors at the same time.
- Identify as many correct terms as possible, while controlling the rate of inclusion of extraneous terms.

19

What is special here?

- Goal: identify the major players
- A continuum of ordinal-valued covariates (the genetic loci)
- Association among the covariates
 - Loci on different chromosomes are independent
 - Along chromosome, a very simple (and known) correlation structure

20

Exploratory methods

- Condition on a large-effect QTL

- Reduce residual variation
- Conditional LOD score:

$$\text{LOD}(q_2 | q_1) = \log_{10} \left\{ \frac{\Pr(\text{data} | q_1, q_2)}{\Pr(\text{data} | q_1)} \right\}$$

- Piece together the putative QTL from the 1d and 2d scans

- Omit loci that no longer look interesting (drop-one-at-a-time analysis)
- Study potential interactions among the identified loci
- Scan for additional loci (perhaps allowing interactions), conditional on these

21

Automation

- Assistance to the masses
- Understanding performance
- Many phenotypes

22

Additive QTL

Simple situation:

- Dense markers
- Complete genotype data
- No epistasis

$$y = \mu + \sum \beta_j q_j + \epsilon \quad \text{which } \beta_j \neq 0?$$

$$p\text{LOD}(\gamma) = \text{LOD}(\gamma) - T |\gamma|$$

$$0 \text{ vs } 1 \text{ QTL: } p\text{LOD}(\emptyset) = 0$$

$$p\text{LOD}(\{\lambda\}) = \text{LOD}(\lambda) - T$$

23

Experience

- Controls rate of inclusion of extraneous terms
- Forward selection over-selects
- Forward selection followed by backward elimination works as well as MCMC
- Need to define performance criteria
- Need large-scale simulations

Epistasis

$$y = \mu + \sum \beta_j q_j + \sum \gamma_{jk} q_j q_k + \epsilon$$

$$p\text{LOD}(\gamma) = \text{LOD}(\gamma) - T_m |\gamma|_m - T_i |\gamma|_i$$

T_m = as chosen previously

T_i = ?

25

Idea 1

Imagine there are two additive QTL and consider a 2d, 2-QTL scan.

$$T_i = 95\text{th percentile of the distribution of} \\ \max \text{LOD}_f(s, t) - \max \text{LOD}_a(s, t)$$

For the mouse genome:

$$T_m = 2.69 \text{ (BC) or } 3.52 \text{ (F}_2\text{)}$$

$$T_i^H = 2.62 \text{ (BC) or } 4.28 \text{ (F}_2\text{)}$$

26

Idea 2

Imagine there is one QTL and consider a 2d, 2-QTL scan.

$$T_m + T_i = 95\text{th percentile of the distribution of} \\ \max \text{LOD}_f(s, t) - \max \text{LOD}_1(s)$$

For the mouse genome:


$$T_m = 2.69 \text{ (BC) or } 3.52 \text{ (F}_2\text{)}$$

$$T_i^H = 2.62 \text{ (BC) or } 4.28 \text{ (F}_2\text{)}$$

$$T_i^L = 1.19 \text{ (BC) or } 2.69 \text{ (F}_2\text{)}$$


27

Models as graphs



28


Results



$$T_m = 2.69 \quad T_i^H = 2.62 \quad T_i^L = 1.19 \quad T_m + T_i^H = 5.31 \quad T_m + T_i^L = 3.88 \quad 2T_m = 5.38$$

29


Add an interaction?



$$T_m = 2.69 \quad T_i^H = 2.62 \quad T_i^L = 1.19 \quad T_m + T_i^H = 5.31 \quad T_m + T_i^L = 3.88 \quad 2T_m = 5.38$$

30


Add another QTL?



$$T_m = 2.69 \quad T_i^H = 2.62 \quad T_i^L = 1.19 \quad T_m + T_i^H = 5.31 \quad T_m + T_i^L = 3.88 \quad 2T_m = 5.38$$

31

Add a pair of QTL?



$$T_m = 2.69 \quad T_i^H = 2.62 \quad T_i^L = 1.19 \quad T_m + T_i^H = 5.31 \quad T_m + T_i^L = 3.88 \quad 2T_m = 5.38$$

32

To do

- Improve search procedures
- Measuring model uncertainty
- Measuring uncertainty in QTL location

33

Summary

- QTL mapping is a model selection problem
- The criterion for comparing models is most important
- We're focusing on a penalized likelihood method and are close to a practiceable solution
- Manichaikul et al., Genetics 181:1077–1086, 2009

34